


All information, content, and material of this website / handout is for informational purposes only and are not intended to serve as a substitute for the consultation, diagnosis, and/or medical treatment of a qualified physician or healthcare provider. The information is not intended to recommend the self-management of health problems or wellness. It is not intended to endorse or recommend any particular type of medical treatment. Should the reader have any health care related questions, that person should promptly call or consult your physician or healthcare provider. This information should not be used by any reader to disregard medical and/or health related advice or provide a basis to delay consultation with a physician or a qualified healthcare provider.

## Pelvic Floor Dysfunction (PFD)

The **pelvic floor** is made up of the bony pelvis (hip bones) together with different layers of muscles, fascia, and ligaments. The pelvic floor acts like a hammock to support the pelvic organs including the uterus, bladder, and rectum. If the muscles become overactive, strained or uncoordinated, they may cause pain in the pelvis. This pain may lead the muscles to not contract, relax, or work together.

### Symptoms

Symptoms related to PFD may include pain of the lower abdomen and pelvic region, a sensation of vaginal heaviness or pressure, pain with vaginal penetration, and low back pain that cannot be explained by other reasons. PFD can also impact bladder and bowel function. Bladder symptoms may include urinary urgency and frequency, feeling of incomplete emptying, intermittent urinary stream or the need to strain, and urinary incontinence (leakage). Bowel symptoms may include constipation, pain with bowel movements, frequent bowel movements and fecal incontinence (leakage of stool.) Symptoms of PFD tend to develop slowly and worsen over time.



### Main causes of PFD

While the cause of PFD is not always known contributing factors may include pregnancy, vaginal delivery, pelvic trauma, pelvic surgery and obesity. PFD is also frequently found alongside pelvic diseases such as endometriosis, bladder pain syndrome, irritable bowel syndrome and vulvar pain. PFD may also arise due to repeated straining (such as with bowel movements) leading to poor coordination of the pelvic floor muscles. The pelvic floor muscles may also be involved in compensating for other musculoskeletal conditions, such as low back or hip pain.

### Treatment

**Physical Therapy** is performed by a physical therapist who has been specifically trained in pelvic health. The physical therapist will perform a complete initial evaluation and, together with the patient, will establish goals and develop an individualized treatment plan. The treatment plan may include patient education, manual therapy, therapeutic exercise, postural training, breathing exercises, neuromuscular reeducation (teaching how to improve pelvic floor muscle control including relaxation, contraction, and coordination), biofeedback, and home exercise program. Modalities such as cold laser, interferential current, electrical stimulation, ultrasound, heat, and ice may also be used.

**Medications** in the form of muscle relaxants or nerve pain medicines can be given to relax the pelvic muscles, desensitize the nervous system, and help the patient tolerate physical therapy.

**Trigger point injections** are injections placed directly in the dysfunctional muscles to control pain, treat inflammation, and reduce spasm. Injections may include a numbing agent, a steroid, or even botulinum toxin.

**PFD** often requires a combination of treatments in addition to physical therapy. In patients with chronic pain, other interventions such as stress control, lifestyle modification, cognitive behavioral therapy (CBT), relationship therapy, meditation, yoga, and acupuncture may be used to reduce pain and improve function.

For more information on Pelvic Floor Dysfunction visit: [www.ichelp.org](http://www.ichelp.org) or [www.fascrs.org](http://www.fascrs.org)